arXiv:0707.4141v2 [math.SG] 9 Dec 2007

RESOLUTION OF SYMPLECTIC CYCLIC ORBIFOLD SINGULARITIES

KLAUS NIEDERKRÜGER AND FEDERICA PASQUOTTO

Abstract. In this paper we present a method to obtain resolutions of symplectic orbifolds arising from symplectic reduction of a Hamiltonian $\mathbb{S}^1$–manifold at a regular value.

As an application, we show that all isolated cyclic singularities of a symplectic orbifold admit a resolution and that pre-quantisations of symplectic orbifolds are symplectically fillable by a smooth manifold.

1. Introduction

Symplectic quotients are an important source of new symplectic manifolds: they appear as *symplectic reductions* in the context of Hamiltonian actions and the associated moment maps. More generally, reduced spaces corresponding to regular values of the moment map turn out to be *symplectic orbifolds*, but one can still look for a closed smooth symplectic manifold which is isomorphic to the orbifold outside a neighbourhood of the singular set: we call this object a *symplectic orbifold resolution*.

Even in the case of a reduced space corresponding to a singular value of the moment map, where singularities of a more complicated type can occur, Kirwan's "partial desingularisation" method [Kir85] can be applied to obtain a resolution which has only orbifold singularities.

The method described in this paper relies on the construction of an auxiliary circle action in a neighbourhood of the singularities with largest structure group: this action is subsequently used to perform a symplectic cut (as described in [Ler95]), which also amounts to a weighted blow up along the singular stratum [God01]. The orbifold obtained in this way has singularities of lower order, and we can repeat the step inductively until we produce a smooth manifold. Making use of this procedure, we are able to find orbifold resolutions for reduced spaces obtained from symplectic reduction at the regular level sets of a Hamiltonian function generating an $\mathbb{S}^1$–action. In particular, we find symplectic resolutions for all isolated cyclic orbifold singularities. A technique involving symplectic resolutions of orbifolds via blow-ups can also be found in [CFM07]: due to a very restrictive definition of orbifolds, though, that result only provides resolutions in the special case of isolated singularities.

As a direct application of our result, we are able to show that a Seifert manifold with a contact structure that is $\mathbb{S}^1$–invariant and transverse to the fibres is symplectically fillable by a smooth manifold.

Acknowledgements. K. Niederkrüger is working at the *ENS de Lyon*, where he is being funded by the Agence Nationale de la Recherche (ANR) project *Symplexe*. F. Pasquotto is working at the *Vrije Universiteit Amsterdam* and is supported by VENI grant 639.031.620 of the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO).

We thank the *Mathematisches Forschungsinstitut Oberwolfach* for inviting us to a *Research in Pairs* stay at the beginning of 2007, during which we developed the main parts of this article.

1.1. **Isolated singularities in dimension four.**

We start our paper by discussing singularities of the simplest form: we hope that this will provide the reader with some motivation and will serve as the right introduction to the difficulties arising when considering more general examples.

In the four-dimensional case, we can give a very explicit description of the resolution of an isolated orbifold singularity. In order to do so, we use weighted blow-ups of isolated symplectic orbifold singularities, as defined in [God01].

Let $x \in \big(M^{(4)}, \omega\big)$ be an isolated orbifold singularity with structure group $\mathbb{Z}_p = \{1, \xi, \xi^2, \dots, \xi^{p-1}\}$. The orbifold chart around x is $\mathbb{Z}_p$–equivariantly symplectomorphic to a neighbourhood of 0 in $\mathbb{C}^2$ with $\mathbb{Z}_p$ acting by

$$(z_1, z_2) \mapsto (\xi^m z_1, \xi z_2), \quad 0 < m < p, \quad \gcd(m, p) = 1 \ .$$

If m and p were not coprime, the orbifold singularity would not be isolated. We can define an $\mathbb{S}^1$–action on $\mathbb{C}^2$ by setting $\lambda \cdot (z_1, z_2) = (\lambda^m z_1, \lambda z_2)$ for $\lambda \in \mathbb{S}^1$. The actions of $\mathbb{Z}_p$ and $\mathbb{S}^1$ commute, but the induced circle action on $\mathbb{C}^2/\mathbb{Z}_p$ is not effective. Instead we have to go to the $\mathbb{S}^1/\mathbb{Z}_p$–action obtained from the following exact sequence

$$0 \longrightarrow \mathbb{Z}_p \longrightarrow \mathbb{S}^1 \longrightarrow \hat{\mathbb{S}}^1 \longrightarrow 0 \ ,$$

with the homomorphism of the circle given by $\lambda \mapsto \lambda^p$. This defines now a symplectic $\hat{\mathbb{S}}^1$–action on $\mathbb{C}^2/\mathbb{Z}_p$ by

$$\mu \cdot [z_1, z_2] = [\lambda \cdot (z_1, z_2)] = [\lambda^m z_1, \lambda z_2]$$

for $\mu \in \hat{\mathbb{S}}^1$ and a $\lambda \in \mathbb{S}^1$ such that $\lambda^p = \mu$.

The weighted blow-up of $\mathbb{C}^2/\mathbb{Z}_p$ at the origin can be represented as a symplectic cut with respect to this $\hat{\mathbb{S}}^1$–action: Take the product manifold $\mathbb{C}^2/\mathbb{Z}_p \times \mathbb{C}$ with the symplectic form given by $(\omega, -i\, dw \wedge d\bar{w})$ and the effective $\hat{\mathbb{S}}^1$–action

$$\mu \cdot \big([z_1, z_2], w\big) = \big(\mu \cdot [z_1, z_2], \mu^{-1} w\big) = \big([\lambda^m z_1, \lambda z_2], \lambda^{-p} w\big), \quad \lambda^p = \mu \ .$$

The blow-up is the symplectic reduction of this space. The Hamiltonian function that corresponds to this action is

$$H\big([z_1, z_2], w\big) = m|z_1|^2 + |z_2|^2 - p|w|^2 \ ,$$

and the ε–level set $H^{-1}(\varepsilon)$ is diffeomorphic to the manifold $\mathbb{S}^3/\mathbb{Z}_p \times \mathbb{C}$. The last step consists in taking the quotient $H^{-1}(\varepsilon)/\hat{\mathbb{S}}^1$ to obtain a symplectic orbifold that can be glued to $\mathbb{C}^2/\mathbb{Z}_p$ after removing a neighbourhood of the origin.

There is only one singular point in $H^{-1}(\varepsilon)/\hat{\mathbb{S}}^1$, namely $\big([1, 0], 0\big) \in \mathbb{S}^3/\mathbb{Z}_p \times \mathbb{C}$, with stabiliser $\mathbb{Z}_m$. By the slice theorem, a neighbourhood of this point admits an orbifold chart equivalent to $\mathbb{C}^2$ with structure group $\mathbb{Z}_m$ acting by $\eta \cdot (w_1, w_2) = (\eta^{-p} w_1, \eta w_2)$.

Now choose $a_1 \in \mathbb{Z}$ such that $0 < a_1 m - p < m$ and set $m_1 = a_1 m - p$ and $p_1 = m$: then the new singularity can also be modelled by $\mathbb{Z}_{p_1}$ acting by $\eta \cdot (z_1, z_2) = (\eta^{m_1} z_1, \eta z_2)$ with $\gcd(m_1, p_1) = 1$, because if b divides both p_1 and m_1 then it also divides m and p. We are thus in the initial type of situation, but we have managed to reduce the order of the singularity.

If we blow up once more, we replace this by a new singularity, this time with structure group $\mathbb{Z}_{m_1}$ acting by $\zeta \cdot (z_1, z_2) = \big(\zeta^{a_2 m_1 - p_1} z_1, \zeta z_2\big)$. If we iterate this blow-up process, at each step we replace the previous singularity by a new one with structure group $\mathbb{Z}_{p_i}$ acting by $(\xi^{m_i} z_1, z_2)$, where the pair (p_i, m_i) is recursively given by

$$\begin{pmatrix} p_i \\ m_i \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & a_i \end{pmatrix} \begin{pmatrix} p_{i-1} \\ m_{i-1} \end{pmatrix} \ ,$$

with each a_i corresponding to the “roundup” of $\frac{p_{i-1}}{m_{i-1}}$, that is, the least integer $\geq \frac{p_{i-1}}{m_{i-1}}$. The sequence $[a_1, a_2, \dots]$ corresponds to the continued fraction of $\frac{p}{m}$, so in particular it is a finite sequence (a description of resolutions in terms of continued fractions is contained, for example, in Miles Reid’s lecture notes [Rei]). After sufficiently many blow-ups, in other words, we get a pair of the form $(p_N, 1)$ and thus an orbifold chart which is in fact smooth. This is then the resolution of our initial singularity.

Notice that we can think of each weighted blow-up as taking the connected sum with a suitable orbifold. In dimension 4, if we define the *weighted projective space* to be

$$\mathbb{CP}(a_0, a_1, a_2) = \Big\{ [z_0 : z_1 : z_2] \sim [\lambda^{a_0} z_0 : \lambda^{a_1} z_1 : \lambda^{a_2} z_2] \ \Big| \ (z_0, z_1, z_2) \in \mathbb{C}^3 - \{0\} \text{ and } \lambda \in \mathbb{C}^* \Big\},$$

the weighted blow-up of a singular point with structure group $\mathbb{Z}_p$ acting by $(z_1, z_2) \mapsto (\xi^a z_1, \xi^b z_2)$ can be described as taking the connected sum, around this point, with the orbifold $\mathbb{CP}(a, b, p)$ with


$\overline{\mathbb{CP}}(1,m,p)$
$\overline{\mathbb{CP}}(1,m_1,p_1)$
$\overline{\mathbb{CP}}(1,1,p_N)$
$M \cong \mathbb{C}^2/\mathbb{Z}_p$

Figure 1. The resolution obtained via a sequence of blow-up can also be thought of as the connected sum $M\#\overline{\mathbb{CP}}(1,m,p)\#\overline{\mathbb{CP}}(1,m_1,p_1)\#\cdots\#\overline{\mathbb{CP}}(1,1,p_N)$.

reversed orientation. We can use this description to represent the resolution of a four-dimensional cyclic singularity as in Fig. 1.

In higher dimension (≥ 6), the method just described is not suitable: even if we start with an isolated singularity, after the first blow-up the singular set is not necessarily discrete any more.

Example 1. Consider an isolated orbifold singularity modelled on a neighbourhood of 0 in $\mathbb{C}^3$ with $\mathbb{Z}_p$ acting by

$$(z_1,z_2,z_3)\mapsto(\xi^{m_1}z_1,\xi^{m_2}z_2,\xi z_3),\quad 0<m_1<m_2<p,\quad \gcd(m_j,p)=1, j=1,2\ .$$

We can blow up this singularity with the method used in the four-dimensional case, that is, performing a symplectic cut with respect to a suitable circle action. If $\gcd(m_1,m_2)=d\neq 1$, though, after blowing up the singular set will be a two-dimensional suborbifold with stabiliser $\mathbb{Z}_d$ at generic points.

One could still hope to obtain a resolution of general cyclic orbifold singularities by using weighted blow-ups along suborbifolds (cf. [MS99]), but for this we would need to find a suitable circle action on the fibres of the normal orbibundle to singular strata. While this is not difficult in individual bundle charts, we were not able to define a global action.

In the rest of this paper we thus adopt a different desingularisation method, which has the advantage of applying to a larger class of orbifold singularities than just cyclic isolated ones.

2. Symplectic orbifolds as quotients of Hamiltonian $\mathbb{S}^1$–manifolds

Let M be a manifold with an action of the unit circle $\mathbb{S}^1$. The infinitesimal action is defined by the vector field

$$X_M(p) := \left.\frac{d}{dt}\right|_{t=0}\exp(tX)*p$$

for every $p\in M$.

Definition. A **Hamiltonian** $\mathbb{S}^1$**–manifold** (M,ω) is a symplectic manifold with an $\mathbb{S}^1$–action for which there exists a Hamiltonian function $H: M\to\mathbb{R}$ that generates the action or, equivalently, that satisfies the identity

$$i_{X_M}\omega = dH\ .$$

If this is the case, the function H is also called the **moment map** of the action.

The following statement is well known ([Wei77]), but for completeness we will briefly sketch the proof.

Proposition 1. *Let* (M,ω) *be a Hamiltonian* $\mathbb{S}^1$*–manifold with Hamiltonian function* H*. For any regular value* E *of* H*, the symplectic quotient*

$$M_E := H^{-1}(E)/\mathbb{S}^1$$

is a symplectic orbifold.

Proof. Since E is a regular value, it follows that $H^{-1}(E)$ is a smooth submanifold on which the $\mathbb{S}^1$–action is semi-free, because $i_{X_M}\omega = dH \neq 0$. The stabilizer of a point $p \in H^{-1}(E)$ is thus isomorphic to some finite cyclic subgroup. By the slice theorem, we can find a homeomorphism from a neighbourhood of the equivalence class of p in the orbit space to the quotient $S_p/\operatorname{Stab}(p)$, where S_p denotes a slice at p (the normal space to the $\mathbb{S}^1$–orbit). In this way we obtain an orbifold atlas for the space M_E. Moreover, the symplectic form ω induces a symplectic structure on the orbifold M_E, because $\mathcal{L}_{X_M}\omega \equiv 0$, and $\omega(X_M,\cdot)|_{TH^{-1}(E)} = dH|_{TH^{-1}(E)} \equiv 0$. □

Example 2. The **weighted projective space** $\mathbb{CP}(a_0, a_1, \dots, a_n)$ is the $2n$–dimensional orbifold obtained as the symplectic quotient of $\mathbb{C}^{n+1}$ by the $\mathbb{S}^1$–action

$$\lambda * (z_0, z_1, \dots, z_n) = (\lambda^{a_0} z_0, \dots, \lambda^{a_n} z_n)\ .$$

It is in fact a symplectic orbifold with the structure induced by the canonical symplectic form on $\mathbb{C}^{n+1}$, namely $\omega_0 = \frac{i}{2}\sum dz_j \wedge d\bar{z}_j$. The Hamiltonian function generating the $\mathbb{S}^1$–action is $H(z_0, \dots, z_n) = a_0|z_0|^2 + \dots + a_n|z_n|^2$.

2.1. Symplectic resolutions.

Definition. Let (M,ω) be a symplectic orbifold with singular set X. A **symplectic ε–resolution** of M consists of a smooth symplectic manifold $(\widetilde{M},\widetilde{\omega})$ and a continuous map $p : (\widetilde{M},\widetilde{\omega}) \to (M,\omega)$ for which there is an orbifold tubular neighbourhood U of X of size ε, such that

$$p|_{\widetilde{M}-p^{-1}(U)} : \widetilde{M} - p^{-1}(U) \to M - U$$

is a symplectic diffeomorphism.

In this paper we prove the following result.

Theorem 2. *Let* M *be a symplectic orbifold arising from symplectic reduction of a Hamiltonian* $\mathbb{S}^1$*–manifold at a regular value of the moment map. Then there exists a symplectic* ε*–resolution of* M *for any arbitrarily small* $\varepsilon > 0$*.*

2.2. **The stratification of the singular set.** Let W be a symplectic $\mathbb{S}^1$–manifold with Hamiltonian H and denote by $\operatorname{Sing} W \subset W$ the singular set of the circle action, that is,

$$\operatorname{Sing} W = \left\{x \in W \;\middle|\; \operatorname{Stab}(x) \neq \{1\}\right\}\ .$$

For a given isotropy group $\mathbb{Z}_k$, let W_k denote the union of orbits whose isotropy group is $\mathbb{Z}_k$, namely $W_k = \{x \in W \mid \operatorname{Stab}(x) \cong \mathbb{Z}_k\}$. Then $\operatorname{Sing} W$ is stratified by singular strata, i.e., connected components of W_k for all $k \neq 1$. Assume that 0 is a regular value of H and denote by P the level set $H^{-1}(0)$ and by $\operatorname{Sing} P$ the singular set of the action on this level set. If $\pi : P \to M = P/\mathbb{S}^1$ denotes the orbit map, $X = \pi(\operatorname{Sing} P)$ is the set of orbifold singularities of M and the stratification of $\operatorname{Sing} P$ descends to a stratification of X. Our desingularisation method works by induction on the order of the stabilisers of the strata W_k. Namely, we start with the stratum with largest stabiliser (*minimal stratum*), remove a small neighbourhood of it and glue in a smooth manifold, in such a way that both the symplectic form and the Hamiltonian $\mathbb{S}^1$–action extend to the manifold resulting from this surgery. The new singular set will also carry a stratification, but the order of the stabiliser of the minimal stratum will be strictly less than k. If we successively repeat this procedure sufficiently often, we will eventually reduce this maximal stabiliser to the trivial group. In particular, symplectic reduction at 0 yields then a smooth symplectic manifold which gives a resolution of the orbifold M.

After working out this desingularisation method for symplectic orbifolds, we were told by one of the authors of [GGK99] that they, and others before them, had already used the same construction in the smooth category.

3. Construction of the resolution

The strategy of our construction is to find an auxilliary circle action on the Hamiltonian $\mathbb{S}^1$–manifold around its minimal stratum. This action will have certain properties (see Proposition 5) that will allow us to perform a symplectic cut [Ler95]. The stabilizers of the Hamiltonian $\mathbb{S}^1$–manifold obtained this way are all lower than the stabilizer of the initial minimal stratum. Then we successively repeat the construction until we obtain a manifold with a free circle action.

3.1. **An auxiliary circle action around the minimal stratum.** Suppose (W, ω) is a Hamiltonian $\mathbb{S}^1$–manifold with Hamiltonian function $H : W \to \mathbb{R}$. Assume further that 0 is a regular level set of H, and choose a metric and an almost complex structure J which are $\mathbb{S}^1$–invariant and compatible with ω (see for example [MS98, Section 5.5]). Since symplectic reduction is a local process, it will be sufficient to study a neighbourhood of $P := H^{-1}(0)$ consisting of regular level sets $P_t := H^{-1}(t)$, $t \in (-\varepsilon, \varepsilon)$. We will start by finding a suitable model for this neighbourhood.

Proposition 3. *There is a neighbourhood of P in (W, ω) that is $\mathbb{S}^1$–diffeomorphic to*

$$P \times (-\varepsilon, \varepsilon) \,,$$

with trivial circle action on the second factor $\lambda * (p, t) = (\lambda * p, t)$. Under this diffeomorphism, the Hamiltonian function pulls back to $H(p, t) = t$.

Proof. Consider the vector field

$$Y := \frac{1}{\|\nabla H\|^2} \nabla H = \frac{1}{\|X_{\mathbb{S}^1}\|^2} J X_{\mathbb{S}^1} \,.$$

It is transverse to the level sets, and allows us to define for small times t with $|t| < \varepsilon$ the diffeomorphism

$$\Psi : \; P \times (-\varepsilon, \varepsilon) \to W, \; (p, t) \mapsto \Phi_t^Y(p)$$

onto a neighbourhood of P in W. The pullback of H gives $H \circ \Psi(p, t) = t$, and the circle action is t–invariant (by which we mean $\lambda * \Psi(p, t) = \Psi(\lambda * p, t)$ for $\lambda \in \mathbb{S}^1$), because Y commutes with the generator of the $\mathbb{S}^1$–action. □

¿From now on, we will always assume the neighbourhood of the level set to be of the form $P \times (-\varepsilon, \varepsilon)$ with an $\mathbb{S}^1$–action on the first factor, and with Hamiltonian function $H(p, t) = t$.

In the next step, we will construct a suitable model for a neighbourhood of the minimal stratum, in which we can describe the auxiliary circle action needed for the resolution.

The $\mathbb{S}^1$–action on W is semi-free. Consider the stratification $\{W_k\}$ of the singular set $\operatorname{Sing} W$ given by the isotropy groups. Each stratum W_k is an $\mathbb{S}^1$–invariant symplectic submanifold of W of the form $P_k \times (-\varepsilon, \varepsilon)$, where $P_k := W_k \cap P$.

If k is maximal, that is, there are no points in W of order larger than k, then $W_k \cap P_t$ is a closed submanifold. By restricting to one component, we may further assume W_k to be connected. Choose any $\mathbb{S}^1$–invariant metric g_P on P and extend it to the product metric $g_W := g_P \oplus dt^2$ on $W = P \times (-\varepsilon, \varepsilon)$. With our choice of metrics on P and W, the corresponding exponential maps are related as follows:

$$\exp^W_{(p,t)}(v + a\,\partial_t) = \big(\exp^P_p(v), t + a\big)$$

with respect to the splitting $T_{(p,t)}W = T_pP \oplus \langle \partial_t \rangle$. Moreover, if we denote the normal bundle of P_k in P by N_k, then the normal bundle of W_k in W is isomorphic to the product $N_k \times (-\varepsilon, \varepsilon)$. Therefore a tubular neighbourhood of W_k in W can also be identified via the exponential map with the product $N_k(\delta) \times (-\varepsilon, \varepsilon)$, where $N_k(\delta)$ denotes the δ–subdiskbundle of N_k. This can be summarized in the following proposition:

Proposition 4. *There exists a neighbourhood of the singular stratum W_k in W that has the form $N_k(\delta) \times (-\varepsilon, \varepsilon)$. The Hamiltonian function on this neighbourhood is just given by $H(v,t) = t$ and the circle action is the linearized $\mathbb{S}^1$–action on N_k.*

In what follows we will therefore implicitly assume this identification and denote for simplicity by ω and J the pullback of the corresponding structures on W.

Later it will be necessary to introduce a second circle action. To avoid confusions, from now on we will call the linearisation of the given action the β–action, and we will write it as $\lambda *_\beta v$ for $\lambda \in \mathbb{S}^1$ and $v \in \nu_k(\delta)$.

Let $x \in W_k$ be a singular point in the minimal stratum. The stabiliser $\operatorname{Stab}(x) \cong \mathbb{Z}_k$ acts by isometric (with respect to both g_W and the metric $g_J := \omega(\cdot, J\cdot)$), J–linear transformations on T_xW, hence the $\mathbb{Z}_k$–action is equivalent to

$$\lambda *_\beta \mathbf{z} = (\lambda^{\widetilde{a}_1} z_1, \dots, \lambda^{\widetilde{a}_n} z_n), \quad \lambda \in \mathbb{Z}_k,\ \mathbf{z} \in T_xW,\ \widetilde{a}_1, \dots, \widetilde{a}_n \in \mathbb{Z}\ .$$

Without loss of generality we may assume that $0 = \widetilde{a}_1 = \dots = \widetilde{a}_m < \widetilde{a}_{m+1} \le \dots \le \widetilde{a}_n < k$ for some m. The first m directions span the space T_xW_k, and the other directions are orthogonal to T_xW_k: it is easy to show that this holds not only for g_J, but also with respect to any other $\mathbb{S}^1$–invariant metric. Hence, if we denote by ν_k the normal bundle of W_k in W, the subspace $\{(0, \dots, 0, z_{m+1}, \dots, z_n)\}$ coincides with the fibre $\nu_k(x)$. For all $x \in W_k$, $j_x := J|_{\nu_k(x)}$ defines a complex vector bundle structure on ν_k, making the $\mathbb{Z}_k$–action j–complex linear in each fibre.

Denote by $a_1 < \dots < a_l$ the *distinct* exponents occurring in the normal form for the action: ν_k splits thus into a direct sum of subbundles

$$\nu_k = E_1 \oplus \dots \oplus E_l\ ,$$

where $E_i(x)$ denotes the eigenspace corresponding to the eigenvalue λ^{a_i} in the fibre at the point x. This splitting is well defined for each component of W_k. Therefore we can extend the $\mathbb{Z}_k$–action to a second circle action by setting for any $\lambda \in \mathbb{S}^1$

$$\lambda *_{\mathrm{aux}} v := \lambda^{a_1} v_1 + \dots + \lambda^{a_l} v_l\ ,$$

where $v = v_1 + \dots + v_l$ is a splitting with respect to the eigenspaces defined above, and $\lambda^{a_i} v_i := \bigl(\cos(a_i\lambda) + \sin(a_i\lambda)\, j\bigr) v_i$. This auxiliary action is fibrewise and j–linear, and commutes with the original β–action. Unfortunately it does not need to respect the symplectic form ω. Recall that ω denotes here the symplectic form on the total space of ν_k obtained by pulling back the symplectic form on W. By averaging ω over the auxiliary action, we obtain a closed 2–form ω' on ν_k that is invariant with respect to both the β– and the auxiliary action. There is a small neighbourhood of the zero section of ν_k where we also have $(\omega')^n \ne 0$ and hence ω' is a symplectic form. To prove this it suffices to show that ω is aux–invariant on the zero section. In fact, at the zero section W_k of ν_k, there is a well defined splitting of the tangent bundle $T\nu_k|_{W_k} = TW_k \oplus \nu_k$ and we can write

$$T\nu_k|_{W_k} = TW_k \oplus E_1 \oplus \dots \oplus E_l\ .$$

The E_i's are J–linear subspaces and g_J–orthogonal to each other, so that they are also symplectically orthogonal. The linearised auxiliary action on a vector $w + v_1 + \dots + v_l \in T\nu_k$ with $w \in T_xW_k$ and $v_i \in E_i(x)$ is given by

$$\lambda *_{\mathrm{aux}} (w + v_1 + \dots + v_l) = w + \lambda^{a_1} v_1 + \dots + \lambda^{a_l} v_l$$

and using the orthogonality relations, it is easy to check that

$$\omega(\lambda *_{\mathrm{aux}} v, \lambda *_{\mathrm{aux}} v') = \omega(v, v')\ .$$

Hence ω is aux–invariant on the zero section of ν_k and we do not change it there by averaging. It follows that there is a small neighbourhood of the zero section, where ω' will be symplectic.

Proposition 5. *There exists a Hamiltonian $\mathbb{S}^1$–action φ in a neighbourhood of W_k with the following properties:*

- (i) *φ leaves W_k pointwise fixed and the stabilizer of points not lying on W_k is a proper subgroup of $\mathbb{Z}_k$;*
- (ii) *φ commutes with the β–action;*

(iii) *the two actions coincide for elements in* $\mathbb{Z}_k \leq \mathbb{S}^1$, *and if* $\lambda *_\varphi x = \sigma *_\beta x$, *then* $\sigma \in \mathbb{Z}_k \leq \mathbb{S}^1$*;*
(iv) *the Hamiltonian function for the* φ*–action is* β*–invariant and constant along* W_k*. Furthermore, it is a Morse–Bott function with critical set* W_k *of index zero.*

Proof. We will obtain the φ–action by a deformation of the existing auxiliary action. By a standard application of the equivariant Moser trick, there exists a β–equivariant isotopy of a neighbourhood of W_k that deforms ω into ω'. The only difficulty comes from the fact that we are dealing with an open set, so that the flow of the Moser vector field X_t does not need to exist up to time 1. But since X_t vanishes on the zero section W_k, after shrinking W_k and the radius of the tubular neighbourhood, we get a smaller set where the isotopy can be defined as the flow of X_t for all $t \in [0, 1]$. The inverse of this isotopy deforms the auxiliary action into the required action φ.

Since the flow of the Moser vector field leaves W_k invariant and is β–equivariant, one can show that properties (i)–(iii) hold for the auxiliary action together with β and this is equivalent to the corresponding statementes for φ and β.

Since ω is φ–invariant, one has that $d i_{X_\varphi}\omega = \mathcal{L}_{X_\varphi}\omega = 0$. For the time being, let U be any tubular neighbourhood of W_k, where φ is defined. The closed 1–form $i_{X_\varphi}\omega$ represents a class in $H^1(U)$ which vanishes if we pull it back to the zero section W_k: Given that $H^1(U) \cong H^1(W_k)$, it follows that $i_{X_\varphi}\omega$ is exact on U, i.e., there exists a function μ_φ such that $i_{X_\varphi}\omega = d\mu_\varphi$. The function μ_φ is uniquely defined up to an additive constant (which we may choose such that $\mu_\varphi \equiv 0$ on W_k) and is β–invariant.

Recall that a Hamiltonian $\mathbb{S}^1$–function is always Morse-Bott (see [MS98, Section 5.5]). The critical set coincides with the set of fixed points of the action, hence in our case with W_k.

The index of a Morse-Bott function is invariant under diffeomorphisms. Therefore in order to compute the index of μ_φ at W_k, it suffices to compute the index of the Hamiltonian function μ_{aux} of the auxiliary action with respect to ω', because μ_φ is the pullback of μ_{aux} under the Moser-flow.

Let ∂_r be the radial vector field on ν_k given by

$$\partial_r(v) = \left.\frac{d}{dt}\right|_{t=1} t \cdot v$$

for $v \in \nu_k$. We will prove that μ_{aux} strictly increases in radial direction, which shows that it has index zero. More precisely, we will prove that $\mathcal{L}_{\partial_r}\mu_{\mathrm{aux}} \geq 0$ with equality only at the zero section. By definition of μ_{aux}, one has $i_{\partial_r} d\mu_{\mathrm{aux}} = \omega'(X_{\mathrm{aux}}, \partial_r)$, so it will suffice to show that there exists a neighbourhood of W_k where $\omega'(X_{\mathrm{aux}}, \partial_r) \geq 0$.

With π denoting the bundle projection $\nu_k \to W_k$, the vertical bundle $V(\nu_k) \leq T\nu_k$ of ν_k can be identified with the pullback bundle

$$\pi^*\nu_k = \big\{(v, w) \in \nu_k \times \nu_k \;\big|\; \pi(v) = \pi(w)\big\} .$$

The identification of $\pi^*(\nu_k)$ and $V(\nu_k)$ goes as follows

$$\chi :\, \pi^*(\nu_k) \to V(\nu_k), \quad (v, w) \mapsto \left.\frac{d}{dt}\right|_{t=0} (v + tw) .$$

Let $v \in \nu_k$, and write it as $v = v_1 + \cdots + v_l$ with respect to the splitting $\nu_k = E_1 \oplus \cdots \oplus E_l$. Then the vectors X_{aux} and ∂_r are given by

$$X_{\mathrm{aux}}(v) = \chi(v, a_1 j v_1 + \cdots + a_l j v_l) \quad \text{and} \quad \partial_r(v) = \chi(v, v)$$

as elements of $V(\nu_k) \cong \pi^*(\nu_k)$. Now assume $v = (x, 0)$ lies in the zero section of ν_k. Then

$$\omega'\big(\chi(v, \textstyle\sum a_i j w_i), \chi(v, w_1 + \cdots + w_l)\big) = \displaystyle\sum_{j=1}^{l} a_j\, \omega'\big(j\chi(v, w_j), \chi(v, w_j)\big) > 0$$

if $w \neq 0$, since the eigenspaces E_j's are ω–orthogonal. By continuity this also holds for all v in a neighbourhood of the zero section and all $w \neq 0$. Hence in particular $\omega'(X_{\mathrm{aux}}, \partial_r) > 0$ on $U - W_k$. □

3.2. **Surgery along the minimal stratum.** In this section we describe how to replace a neighbourhood of (one component of) the minimal stratum W_k by a smooth manifold, and extend both the symplectic form ω and the Hamiltonian $\mathbb{S}^1$–action β to the resulting manifold, in such a way that the singular points of the extended action are of order strictly smaller than k. In order to achieve this, we perform a symplectic cut with respect to the Hamiltonian $\mathbb{S}^1$–action φ constructed in the previous section. The main reference for symplectic cuts and hence for everything that follows is Lerman's original paper [Ler95].

As in the previous section, consider the minimal singular stratum W_k, denote by U a tubular neighbourhood in W, where the φ–action is defined, and take now the product $U \times \mathbb{C}$. It admits a first circle action, which is just the extension of the original $\mathbb{S}^1$–action β by the trivial action on the $\mathbb{C}$–factor, namely, for $x \in U$, $z \in \mathbb{C}$

$$\lambda *_\beta (x, z) := (\lambda *_\beta x, z) ,$$

and we can define a second circle action on $U \times \mathbb{C}$ by setting

$$\lambda *_\varphi (x, z) = (\lambda *_\varphi x, \lambda^{-k} z) .$$

These two actions commute and therefore we can combine them and define a new $\mathbb{S}^1$–action

$$\lambda *_\tau (x, z) := \lambda *_\varphi \left(\lambda^{-1} *_\beta (x, z)\right) .$$

This τ–action is not effective, because the φ– and the β–action coincide for elements in $\mathbb{Z}_k$. Hence consider the short exact sequence

$$0 \to \mathbb{Z}_k \to \mathbb{S}^1 \to \hat{\mathbb{S}}^1 \to 0 ,$$

with the homomorphism of the circle given by $\lambda \mapsto \lambda^k$, and let $\hat{\mathbb{S}}^1$ act on $U \times \mathbb{C}$ by $\sigma *_{\hat\tau} (x, z) = \lambda *_\tau (x, z)$ for some $\lambda \in \mathbb{S}^1$ such that $\lambda^k = \sigma$. This new action, which we denote by $\hat\tau$, is not only effective but by Proposition 5.(iii), even free and the quotient $(U \times \mathbb{C})/\hat\tau$ is a smooth manifold. It still carries an $\mathbb{S}^1$–action induced by the φ–action on $U \times \mathbb{C}$, and this is well defined because φ commutes with $\hat\tau$.

We define a symplectic form $\Omega = (\omega, -i\, dz \wedge d\bar z)$ on $U \times \mathbb{C}$, which is invariant with respect to the $\hat\tau$–action. By construction, the infinitesimal generator of this action can be written as $X_{\hat\tau} = -X_\beta + X_\varphi$. The Hamiltonian for the $\hat\tau$–action is given by

$$H_{\hat\tau}(x, z) = \mu_\varphi(x) - k|z|^2 - H(x) ,$$

and if we now do symplectic reduction at some level ε we get the quotient $H_{\hat\tau}^{-1}(\varepsilon)/\hat\tau$ which, with the structure induced by Ω and φ, is a smooth Hamiltonian $\mathbb{S}^1$–manifold with Hamiltonian function $H_\varphi[(x, z)] = \mu_\varphi(x) - k|z|^2 = H(x) + \varepsilon$. Notice that $H_{\hat\tau}^{-1}(\varepsilon)$ can be written as the disjoint union of two $\hat\tau$–invariant manifolds

$$H_{\hat\tau}^{-1}(\varepsilon) = \left\{(x, z) \,\middle|\, \mu_\varphi(x) - H(x) > \varepsilon,\ |z|^2 = \frac{\mu_\varphi(x) - H(x) - \varepsilon}{k}\right\} \sqcup \left\{(x, 0) \,\middle|\, \mu_\varphi(x) - H(x) = \varepsilon\right\} .$$

Since the level sets of μ_φ and H are transverse, the restriction of μ_φ to $P = H^{-1}(0)$ is still a Morse-Bott function of index zero. Hence we can choose $\delta > 0$ such that $\mu_\varphi^{-1}(\delta) \cap P$ is contained in the interior of U and has the structure of a sphere bundle over $P_k = P \cap W_k$. Choose $\varepsilon' > 0$ such that

$$\left(\mu_\varphi - H\right)^{-1}(\delta/2, \delta) \cap P_t$$

is non-empty, does not intersect W_k, and is contained in the interior of U for all $|t| < \varepsilon'$ (see Fig. 2). Assume from now on that all subsets in W are restricted to $H^{-1}\big((-\varepsilon', \varepsilon')\big)$. Then

$$U(\delta/2, \delta) := \left(\mu_\varphi - H\right)^{-1}(\delta/2, \delta)$$

is diffeomorphic to a spherical shell bundle over W_k. Choose now $\varepsilon = \delta/2$, set $V := H_{\hat\tau}^{-1}(\varepsilon)/\hat\tau$, and consider the map

$$\Phi : U(\varepsilon, 2\varepsilon) \to V, \qquad x \mapsto \left[x, \sqrt{\frac{\mu_\varphi(x) - H(x) - \varepsilon}{k}}\right] .$$

This is a diffeomorphism (onto its image), equivariant with respect to the β–action on $U(\varepsilon, 2\varepsilon)$ and the φ–action on V. Its inverse can be constructed as follows: given $[x, z]$ with $z \neq 0$, we first represent the same class by an element (x', z') such that z' is a real positive number, and then define

$$\Phi^{-1}([x, z]) := x' \ .$$

Moreover, since Φ factors through a map $U(\varepsilon, 2\varepsilon) \to H_{\hat{\tau}}^{-1}(\varepsilon)$ which is the identity in the first component and a real function in the second one, we have

$$\Phi^*(\omega, -i\, dz \wedge d\bar{z}) = \omega \ ,$$

hence Φ gives in fact an equivariant symplectic identification of $U(\varepsilon, 2\varepsilon)$ with its image under Φ. More precisely we have

$$\Phi\big(U(\varepsilon, 2\varepsilon)\big) = \big\{[x, z] \in V \ \big| \ \varepsilon < \mu_\varphi(x) - H(x) < 2\varepsilon\big\} \ .$$

We can now remove a tubular ε–neighbourhood $\overline{U(\varepsilon)}$ of W_k in W and glue in the smooth manifold

$$V(\varepsilon) := \left\{(x, z) \ \middle| \ \varepsilon \leq \mu_\varphi(x) - H(x) < 2\varepsilon, \ |z|^2 = \frac{\mu_\varphi(x) - H(x) - \varepsilon}{k}\right\} \Big/ \ \hat{\tau}$$

along the open "collar" $U(\varepsilon, 2\varepsilon)$, using the map Φ. In this way we define the new manifold

$$\widetilde{W} = \big(W - \overline{U(\varepsilon)}\big) \cup_\Phi V(\varepsilon) \ .$$

Since Φ is equivariant, the β–action on $W - \overline{U(\varepsilon)}$ and the φ–action on $V(\varepsilon)$ fit together to give a circle action $\widetilde{\beta}$ on $\widetilde{W}$, which by construction coincides with β outside a 2ε–neighbourhood of W_k. Moreover, Φ identifies the given symplectic forms on the two sides of the gluing, so $\widetilde{W}$ also admits a symplectic form $\widetilde{\omega}$ with the property that $\widetilde{\omega} = \omega$ on $W - U(2\varepsilon)$. With the action $\widetilde{\beta}$ and the symplectic form $\widetilde{\omega}$ just defined, $\widetilde{W}$ is a Hamiltonian $\mathbb{S}^1$–manifold. The Hamiltonian function $\widetilde{H}$ for $\widetilde{\beta}$ is given by H on $W - U(2\varepsilon)$ and by $H_\varphi - \varepsilon$ on $V(\varepsilon)$.


V_k
V
Φ
W_k
Φ
$H^{-1}(0) = P$
$H^{-1}\big[(\varepsilon', \varepsilon')\big]$
$U(\varepsilon, 2\varepsilon)$

FIGURE 2. Remove the $U(\varepsilon)$–neighbourhood of W_k, and glue in the $V(\varepsilon)$–patch, identifying along $U(\varepsilon, 2\varepsilon)$ via the diffeomorphism Φ.

We need to analyse the singular points of the $\widetilde{\beta}$–action on the "patch" $V(\varepsilon)$. They satisfy the relation $\lambda *_\varphi [x, z] = [x, z]$ for some $\lambda \in \mathbb{S}^1$ and this in turn means that there exist $\kappa \in \hat{\mathbb{S}}^1$ and $\sigma \in \mathbb{S}^1$, $\sigma^k = \kappa$ such that

$$\lambda *_\varphi (x, z) = \kappa *_{\hat{\tau}} (x, z) = \sigma *_\varphi \sigma^{-1} *_\beta (x, z) \ .$$

In particular, $\lambda *_\varphi x = \sigma *_\varphi \sigma^{-1} *_\beta x$, and by Proposition 5.(iii) this identity can only hold if $\sigma \in \mathbb{Z}_k$. Hence $\kappa = 1$ and singular points are characterised by $\lambda *_\varphi (x, z) = (x, z)$. Since $[x, z] \in V(\varepsilon)$ implies that x does not lie in W_k, the isotropy groups are proper subgroups of $\mathbb{Z}_k$, see Proposition 5.(i).

We have thus proved:

Proposition 6. *Let W_k be the minimal singular stratum of a Hamiltonian $\mathbb{S}^1$–manifold W. There exists a new symplectic manifold $\widetilde{W}$ which is symplectomorphic to W outside an arbitrarily small neighbourhood of W_k and admits a Hamiltonian $\mathbb{S}^1$–action that coincides with the given action away from W_k and only has singularities of order strictly smaller than k.*

In fact we have also shown that 0 is still a regular value of the moment map of the circle action on $\widetilde{W}$ and if we consider the symplectic reduced space at this level, namely

$$\widetilde{M} := \widetilde{H}^{-1}(0)/\widetilde{\beta}\ ,$$

we see that this symplectic orbifold has singularities of strictly lower order than those of $M = H^{-1}(0)/\beta$.

Moreover, there exists a map $f : \widetilde{M} \to M$, which is a symplectic orbifold isomorphism outside an arbitrarily small neighbourhood of $M_k = P_k/\beta$ (and in fact coincides with the identity map outside a slightly larger neighbourhood). We shall describe how to define f. On $(P - U(2\varepsilon))/\beta$ it is simply the identity. In order to define it on $\big(V(\varepsilon) \cap H_\varphi^{-1}(\varepsilon)\big)/\varphi$ a little more work is required. First of all, denote by V_k the quotient

$$\big\{(x, 0) \in H_{\hat{\tau}}^{-1}(\varepsilon) \;\big|\; \mu_\varphi(x) - H(x) = \varepsilon\big\}/\hat{\tau}\ .$$

Then the inverse of the gluing map Φ restricts to a diffeomorphism $\Phi^{-1} : (V - V_k) \cap H_\varphi^{-1}(\varepsilon) \to P \cap U(\varepsilon, 2\varepsilon)$. Since Φ is equivariant with respect to the φ– and β–actions, this descends to a symplectic orbifold isomorphism on the quotients. Let $h : U(\varepsilon, 2\varepsilon) \cap P \to (U(2\varepsilon) - W_k) \cap P$ be a β– and φ–equivariant diffeomorphism (recall that φ is the action on W constructed in Section 3.1), that is the identity in a neighbourhood of the outer boundary of $U(2\varepsilon)$ and extends to a smooth map from $U(2\varepsilon) \cap P$ to itself which maps $U(\varepsilon) \cap P$ to P_k.

Then we can define

$$\begin{aligned} f :\quad & \big(V \cap H_\varphi^{-1}(\varepsilon)\big)/\varphi \to \big(P \cap U(2\varepsilon)\big)/\beta \\ & [x, z] \mod \varphi \mapsto \begin{cases} h \circ \Phi^{-1}([x, z]) \mod \beta & \text{if } z \neq 0, \\ h(x) \mod \beta & \text{if } z = 0. \end{cases} \end{aligned}$$

Because of the boundary conditions on h, the map f extends on the outer side to the identity map. To see that f is continuous in a neighbourhood of $\big(V_k \cap H_\varphi^{-1}(\varepsilon)\big)/\varphi$, one has to show that for any sequence $[x_k, z_k]/\varphi \subset \big(V \cap H_\varphi^{-1}(\varepsilon)\big)/\varphi$ that converges to some element $[x_0, 0]/\varphi$, it follows that $f\big([x_k, z_k]/\varphi\big)$ converges to $f\big([x_0, 0]/\varphi\big)$. We can find representatives (x_k', z_k') for the sequence that converge to $(x_0', 0)$ and such that z_k' is a non-negative real number, and hence $f\big([x_k', z_k']/\varphi\big) = h(x_k')/\beta$ converges to $h(x_0')/\beta = f([x_0', 0]/\varphi) = f([x_0, 0]/\varphi)$.

3.3. **Proof of Theorem 2.** If we inductively repeat the above surgery procedure along the minimal stratum, we will eventually obtain a symplectic manifold $\widetilde{W}$ with a free Hamiltonian $\mathbb{S}^1$–action $\widetilde{\beta}$. The symplectic reduced space of this action at the level 0 will be a smooth symplectic manifold $\widetilde{M}$ and it will be equipped with a function $f : \widetilde{M} \to M$, which factors through all the previous steps of the resolution (that is, reduced spaces with orbifold singularities of decreasing order), and is a symplectic diffeomorphism outside an arbitrarily small neighbourhood of the set of orbifold singularities of M.

4. Applications

4.1. **Isolated cyclic orbifold singularities.** Any isolated cyclic orbifold singularity can be represented as $\mathbb{C}^n/\mathbb{Z}_k$ with symplectic form $\omega = i\sum_{j=1}^n dz_j \wedge d\bar{z}_j$, where the generator $\xi = e^{2\pi i/k}$ of $\mathbb{Z}_k$ acts by

$$\xi \cdot \mathbf{z} = \big(e^{2\pi i/k} z_1, e^{2\pi i a_2/k} z_2, \dots, e^{2\pi i a_n/k} z_n\big)\ ,$$

and $a_2, \dots, a_n \in \mathbb{N}$ are all coprime with k.

To find a resolution of the singularity using Theorem 2, just note that $\mathbb{C}^n/\mathbb{Z}_k$ can be obtained by doing symplectic reduction on the manifold $\mathbb{C}^* \times \mathbb{C}^n$ with $\mathbb{S}^1$–action given by

$$e^{i\varphi} \cdot \big(\rho e^{i\vartheta}, z_1, \dots, z_n\big) := \big(\rho e^{i(k\varphi + \vartheta)}, e^{i\varphi} z_1, e^{i a_2 \varphi} z_2, \dots, e^{i a_n \varphi} z_n\big)\ ,$$

symplectic form $\omega + \rho\, d\rho \wedge d\vartheta + \frac{1}{k}\, d\vartheta \wedge \sum_{j=1}^{n} a_j\, d|z_j|^2$ and moment map $H(\rho e^{i\vartheta}, z_1, \dots, z_n) = \rho^2$. Theorem 2 then immediately implies the following generalisation of the desingularisation result obtained in dimension 4:

Corollary 7. *Every symplectic orbifold with only isolated cyclic singularities admits a symplectic resolution.*

Cavalcanti, Fernández, and Muñoz [CFM07] recently introduced a method that allows one to find resolutions for all isolated symplectic orbifold singularities.

4.2. Generalised Boothby-Wang fibrations are fillable.

Definition. A **Boothby-Wang fibration** is a closed contact manifold (P, α) with a free $\mathbb{S}^1$–action which is given by the flow of the Reeb field X_{Reeb}. A **generalised Boothby-Wang fibration** is a closed contact manifold (P, α), where the Reeb field induces a semi-free $\mathbb{S}^1$–action.

Remark 1. A Boothby-Wang fibration (P, α) defines an $\mathbb{S}^1$–principal bundle over the manifold $B = P/\mathbb{S}^1$ with connection 1–form α. The curvature form ω is the unique 2–form on B that satisfies $\pi^*\omega = d\alpha$. The base manifold (B, ω) is a symplectic manifold and ω represents an integral cohomology class.

Conversely, for any symplectic manifold (B, ω) with integral symplectic form, one can construct a Boothby-Wang fibration (P, α) over it, the so-called **pre-quantisation**. This is the inverse of the previous construction.

Remark 2. A generalised Boothby-Wang fibration can be considered as the pre-quantisation of the symplectic *orbifold* $(P/\mathbb{S}^1, \omega)$, and all the statements made in Remark 1 can be translated to this setting.

Proposition 8. *A generalised Boothby-Wang fibration (P, α) has a natural convex filling by a symplectic orbifold.*

Proof. These computations were obtained with the help of H. Geiges. Consider the (complex) "line bundle" L associated to P, i.e. the bundle obtained from $P \times \mathbb{C}$ by identifying (p, z) with $(e^{-i\varphi} * p, e^{i\varphi}\, z)$ for every $e^{i\varphi} \in \mathbb{S}^1$. The manifold P embeds naturally via

$$P \hookrightarrow L, \quad p \mapsto [p, 1] .$$

The two forms

$$\frac{1}{2}\left(|z|^2\, \alpha + x\, dy - y\, dx\right) \qquad \text{and} \qquad \frac{1}{2}\, d\alpha$$

on $P \times \mathbb{C}$ induce well-defined forms on L. By adding the differential of the first form to the second one, we obtain a pre-symplectic form

$$\omega := \frac{1}{2}\, d(|z|^2) \wedge \alpha + dx \wedge dy + \frac{1 + |z|^2}{2}\, d\alpha ,$$

because $2^n\, \omega^n = n\, (1 + |z|^2)^{n-1}\, (d\alpha)^{n-1} \wedge \left(d|z|^2 \wedge \alpha + 2dx \wedge dy\right)$ has only a one-dimensional kernel on $P \times \mathbb{C}$ that is generated by $-Z_P + x\, \partial_y - y\, \partial_x$. It follows that $(P \times \mathbb{C}, \omega)$ is a pre-symplectic $\mathbb{S}^1$–manifold, and hence L is a symplectic orbifold where all orbifold singularities sit along the zero section.

Finally, the following field

$$X := \frac{1 + r^2}{2r}\, \partial_r = \frac{1 + x^2 + y^2}{2\,(x^2 + y^2)}\, (x\, \partial_x + y\, \partial_y)$$

is a Liouville vector field for the manifold (P, α), and (L, ω) is hence a convex filling of P. □

All the orbifold singularities lie in the interior of L. By passing to a symplectic resolution of L whose existence is guaranteed by Theorem 2, we can obtain a smooth symplectic filling.

Corollary 9. *Generalised Boothby-Wang fibrations are symplectically fillable by a* smooth *manifold.*

Remark 3.
1. Popescu-Pampu recently proved the following conjecture of Biran: there exist prequantizations (for example on higher dimensional tori) that are not holomorphically fillable [PP07]. These manifolds, on the other hand, do have a strong symplectic filling, showing that the different types of fillability do not coincide (a result well-known in dimension 3).
2. Massot showed that any contact structure on a 3–dimensional Seifert manifold that is transverse to the fibres (but not necessarily invariant) has a weak symplectic filling [Mas07]. To achieve this result he constructs first a filling by an orbifold, whose singularities can then be resolved by using for example the method presented in this paper.

References

[CFM07] G. Cavalcanti, M. Fernández, and V. Muñoz, *Symplectic resolutions, lefschetz property and formality*, arXiv **math.SG/arXiv:0710.0731** (2007).

[GGK99] V. Ginzburg, V. Guillemin, and Y. Karshon, *The relation between compact and non-compact equivariant cobordisms*, Tel Aviv Topology Conference: Rothenberg Festschrift (1998), Contemp. Math., vol. 231, Amer. Math. Soc., Providence, RI, 1999, pp. 99–112.

[God01] L. Godinho, *Blowing up symplectic orbifolds*, Ann. Global Anal. Geom. **20** (2001), no. 2, 117–162.

[Kir85] F. Kirwan, *Partial desingularisations of quotients of nonsingular varieties and their Betti numbers*, Ann. of Math. (2) **122** (1985), no. 1, 41–85.

[Ler95] E. Lerman, *Symplectic cuts*, Math. Res. Lett. **2** (1995), no. 3, 247–258.

[Mas07] P. Massot, *Geodesible contact structures on* 3*–manifolds*, arXiv **math/0711.0377** (2007).

[MS98] D. McDuff and D. Salamon, *Introduction to symplectic topology. 2nd ed.*, Oxford Mathematical Monographs. New York, NY: Oxford University Press., 1998.

[MS99] E. Meinrenken and R. Sjamaar, *Singular reduction and quantization*, Topology **38** (1999), no. 4, 699–762.

[PP07] P. Popescu-Pampu, *Holomorphic fillability and cohomology*, arXiv **math/0711.2941** (2007).

[Rei] M. Reid, *Cyclic surface quotient singularities*, http://www.maths.warwick.ac.uk/∼miles/surf/more/cyclic.pdf.

[Wei77] A. Weinstein, *Symplectic V–manifolds, periodic orbits of Hamiltonian systems, and the volume of certain Riemannian manifolds*, Comm. Pure Appl. Math. **30** (1977), no. 2, 265–271.

E-mail address, K. Niederkrüger: `kniederk@umpa.ens-lyon.fr`
E-mail address, F. Pasquotto: `pasquott@few.vu.nl`

(K. Niederkrüger) École Normale Supérieure de Lyon, Unité de Mathématiques Pures et Appliquées, UMR CNRS 5669, 46, allée d'Italie, 69364 LYON Cedex 07, France

(F. Pasquotto) Department of Mathematics, Faculty of Sciences, Vrije Universiteit, De Boelelaan 1081a, 1081 HV Amsterdam, The Netherlands